

Urinary Tract Pathogens: Analysis and Antimicrobial Effects of *Ocimum Gratissimum* Leaf

Favour Obioma Barnabas¹, Ponchang Wuyep², Suzan Paulinus Ukpa¹, Eno Chongs Mantu¹, Uchejeso Mark Obeta^{3*}, Suzan Joseph Nduke¹ and Uju Uchenna Ashien⁴

¹Department of Medical Microbiology, Federal School of Medical Laboratory Science, Jos, Plateau State Nigeria.

²Department of Plant Science and Technology, Faculty of Applied Natural Sciences, University of Jos, Plateau State Nigeria.

³Department of Medical Laboratory Management, Federal School of Medical Laboratory Science, Jos, Plateau State Nigeria.

⁴Department of Basic Sciences, Federal School of Medical Laboratory Science, Jos, Plateau State Nigeria.

*Correspondence author

Uchejeso Mark Obeta

Department of Medical Laboratory Management
Federal School of Medical Laboratory Science
Jos
Plateau State
Nigeria

Submitted : 22 Aug 2022 ; Published : 19 Sept 2022

Citation: Uchejeso Mark Obeta, Urinary Tract Pathogens: Analysis and Antimicrobial Effects Of *Ocimum gratissimum* Leaf . J N food sci tech, 2022; 3(2):1-13.

Abstract

The search for alternative medicine for the treatment of pathogenic organisms associated with urinary tract infection (UTI) is on increase. The study was carried out to determine the antimicrobial activities of *Ocimum gratissimum* leaf extracts on three urinary tract pathogens; *Escherichia coli*, *Staphylococcus aureus* and *Candida albicans*, using their corresponding standard organisms as controls. The phytochemical analysis of aqueous and alcoholic contents of *Ocimum gratissimum* leaf extracts revealed the presence of; alkaloids, carbohydrates, flavonoids, glycosides, phenolics, saponins, steroids, and tannins. The minimum inhibitory concentration (MIC), minimum bacteriostatic concentration (MBC) and minimum fungistatic concentration (MFC) of the extracts tested on the three uropathogens was 62.5 to 125 mg/ml. The ethanolic extracts at the concentration of 500mg/ml exhibited strong activity against the urinary pathogens with maximum zones of inhibitions of 23.5±1.5, 23.6±0.6 and 18.25±1.3mm while the aqueous extracts at the same concentration showed weak antimicrobial activity at 3.5±0.5mm as the highest zone of inhibition and 1.4±0.0mm as the lowest zone of inhibition. Notably, the inhibition zones increased with increase in extracts concentration and decreased with decrease in the concentration of extracts. Meanwhile, 12 fractions were eluted from *O. gratissimum* leave extract. The major phytochemical fractions were mainly alkaloids, flavonoid, and saponins. This study showed that the ethanolic extracts of *Ocimum gratissimum* is a good antimicrobial agent in-vitro and should not be neglected as an alternative medicine for treatment of pathogenic UTIs.

Keywords : UTI, Uropathogens, *Ocimum gratissimum*, Antimicrobial, Alternative medicine.

Introduction

The use of medicinal plants as traditional and alternative medicine is well known in rural areas of many developing countries and has been globally accepted and encouraged (WHO, 2016; Ezekwesili-Ofili & Okaka, 2019; Obeta et al., 2020). In Nigeria, *Ocimum gratissimum* (scent leave) have been used for making soup and stew (Obeta et al., 2021), eradication of Mosquitoes (Agbalaka et al., 2021; Anzaku et al., 2021) and for various medicinal uses (Bhavani et al., 2021; Lunyera et al., 2016). Nwinyi et al. (2009) investigated the antibacterial activity of the aqueous extracts of *Ocimum gratissimum* against *Escherichia coli* and *Staphylococcus aureus*. Dhanalakshmi et al. (2013) and Chijioke et al. (2014) also investigated the antibacterial activity of some medicinal plants used against Uropathogen pathogens and concluded that extracts of plants origin has remarkable anti microbial activity as compared

to antibiotic activity. It was made known that organisms are gaining resistance day by day towards most of the anti biotics used for the treatment of UTI. As a result of this, so many natural products are being tried to overcome the antibiotic resistant action of these organisms. The present study has been undertaken to identify effective herbal medicines to control UTI since herbal medicines are available in our environment and believed to be safe when compared to antibiotics. If these medicinal plants continue to show antimicrobial activities against most pathogenic organisms that threatens human health, it will be an evidence that natural medicine can complement or take the place of antibiotics in future (Dhanalakshmi, 2013; Bazzaz et al., 2021) .

It has also been investigated that the essential oil of *Ocimum*

gratissimum contains eugenol and shows some evidence of antibacterial activity (Nweze & Eze, 2009). A polyherbal preparation of aqueous extract obtained from the leaves of *Gongronema latifolia*, *Vernonia amygdalina* and *Ocimum gratissimum* showed analgesic activity against many pathogenic organisms (Iroanya et al., 2009). The essential oil has potential for use as a food preservative (Nguefack et al., 2009) and is toxic to *Leishmania*. The oil of *Ocimum gratissimum* has been formulated into creams for clinical trials where favourable against certain dermatological disorders caused by pathogenic microbes (Sofowora, 1993). Many scientists have carried out researches on *Ocimum gratissimum* leaf extracts and discovered that the leaf possess antidiabetic properties and anti-hyperlipidemic effect against many pathogenic organisms (Owoyele et al., 2005), (Egesie et al., 2006), (Ayinla et al., 2011), (Shittu et al., 2016).

Lexa et al(2006) was able to demonstrate the anti fungal property of *Ocimum gratissimum* and discovered that the chloroformic fraction of the extract inhibited 23 isolates (92%) of *Cryptococcus neoformans* at a concentration of 62.5µg/ml (Silva et al., 2007) in vitro against human pathogenic dermatophytes (Lexa et al., 2006). The essential oil of *O. gratissimum* has antimicrobial activities against pathogenic strains of Gram positive (*S. aureus*, *Bacillus* spp) and Gram negative (*E. coli*, *P. aeruginosae*, *S. typhi*, *K. pneumonia*, *P.mirabilis*) bacteria and pathogenic fungus *C. albicans* (Matasyoh et al., 2007). *Ocimum gratissimum* is rich in alkaloid, tannins, phytates, flavonoids, oligosaccharides and has tolerable cyanogenic content (Nakaruma et al., 1999), (Lemos et al., 2005). The plant contains terpenoids, eugenol, thymol, saponins and alkaloids. The essential oil of *O. gratissimum* has antimicrobial activities against pathogenic strains of Gram positive (*S. aureus*, *Bacillus* sp) and Gram negative (*E. coli*, *P. aeruginosae*, *S. typhi*, *K. pneumonia*, *P.mirabilis*) bacteria and pathogenic fungus *C. albicans* (Matasyoh et al., 2007).

In traditional medicine, the leaves have been used in the treatment of UTIs, rubbed between palms to prevent cold, sniffed inside the nose to clear blocked nostrils, used as anti-diarrhea agent, and for the treatment of conjunctivitis by instilling directly into the eyes. Despite all these, there is no evidence to back it up.

This study was carried out to evaluate the antimicrobial effects of ethanolic and aqueous extracts of *Ocimum gratissimum* L. (Lamiaceae) used in traditional medicine for the treatment of several ailments such as urinary tract infection on clinical isolates namely: *Escherichia coli* and *Staphylococcus aureus* and *Candida albicans* in comparison with a typed bacterium of *E.coli* (ATCC 11775), *S. aureus* (ATCC 6538) and typed fungal strain of *Candida albicans* (ATCC 10231).

The specific objectives was to carry out *Ocimum gratissimum* leaf extraction; to determine the in vitro activities on *E. coli*, *S. aureus* and *C. albicans* respectively through antimicrobial susceptibility assay; to determine the Minimum Inhibitory Concentration (MIC) and Minimum Bacteriocidal

Concentration (MBC) and to identify the bioactive compounds and fractions in the extracts that has the highest antimicrobial activity against the UTI isolates.

Materials and Methods

Collection and Preparation of *Ocimum gratissimum* leaf

The *Ocimum gratissimum* leaves as shown in figure 1 used for this study was collected from Polo area of Jos in Jos North LGA, Plateau State within the month of July, 2020. It was identified in Herbarium of Plant Science Department in University of Jos with voucher number UJH16000270. The leaves were washed and allowed to air dry and then grinded into fine powder in line with the method described by Nweze et al. (2009). The extract was sieved, labeled, weighed into sterile dry containers and stored at room temperature for use. The two solvents used for the extraction of the leaves were Water, and Alcohol.



Figure 1: *Ocimum gratissimum* Leaf

Extraction Method

Aqueous extraction was carried out using the cold maceration method according to the method described by Ncube et al. (2008) while the method described by Nweze et al. (2009) was used for the ethanolic extractions. The extracts were concentrated at reduced pressure to dryness at 40°C using rotary evaporator and a greenish black coloured sticky residue was obtained which was used to calculate the yield of each extract (formula 1). The extracts were then stored in refrigerator at 4°C.

$$\text{Percentage yield of extracts} = \frac{W_2 - W_1}{W_0} \times 100 \quad (1)$$

Where:

W₂ is the weight of the extract and the container,

W₁ is the weight of the container alone and

W₀ is the weight of the initial dried sample

Phytochemical Analysis of Extracts

Phytochemical analysis of each extract was carried out using the description of Kolkate et al. (2003) to determine the presence or absence of phenols, alkaloids, flavonoids, tannins, saponins, steroids and glycosides. Identification of Fractions was carried out using Thin-Layer chromatography (TLC) and column chromatography respectively.

The Preparation of the TLC Plates, choice of Solvent System (hexane, ethyl acetate and water (4:2:1), preparation of stock extracts to be spotted, development of Thin layer chromatography (TLC), procedure for column chromatography, detection of bands and calculation of retardation factor (Rf) was done using formular (2) according to method earlier described by Harwood et al. (1980) and Martins (2013).

$$Rf = \frac{\text{Distance traveled by the solute}}{\text{Distance traveled by the solvent front TLC plate}} \quad (2)$$

Antimicrobial activity of the fractions was carried out on the 5 fractions of the leaves against the 3 different organisms used. The zones of inhibitions were measured and recorded.

Collection of Standard Strain

Standard strains species which were used as controls were; *Escherichia coli* (ATCC 11775), *Staphylococcus aureus* (ATCC 6538) and *Candida albicans* (ATCC 10231) obtained from Microbiology laboratory of National Veterinary Research Institute (NVRI) Vom. The *Staphylococcus aureus* strains was cultured on Blood agar plates at 37°C and maintained on nutrient agar slants while the *Candida albicans* strain was grown on Sabroud dextrose agar (SDA) plate and maintained on SDA slant at 4°C.

Collection and culturing of urine samples

Already confirmed urine samples of patients suffering from urinary tract infection were collected from the Laboratory of Bingham University Teaching Hospital Jos.

Identification of organisms

Colonies from the cultured plates were selected and characterized on the basis of morphological, cultural, physiological and biochemical characteristics.

Standardization of Test Organisms

Standardization of test organisms was done in Microbiology laboratory of Jos University. The suspension was incubated at 37°C for 24hours and the size was adjusted by diluting each suspension with sterile broth to the 0.5 MacFarland standard turbidity (CLSI, 1998) approximately 1.5×10^6 CFU/ml.

Antimicrobial susceptibility testing

The antimicrobial activities of the extracts were tested on the organisms using the agar-well diffusion method according to method described by Pottumarthy et al., 2006 using Mueller-Hinton agar plates.

Assessment of Minimum Inhibitory Concentration (MIC)

The minimum inhibitory concentration was determined according to national committee for clinical laboratory standard (NCCLS) (1998) using standard two-fold dilution broth methodology.

Determination of Minimum Bacteriocidal Concentration and Minimum Fungicidal Concentration (MBC and MFC)

5ul of each extract from each tube that exhibited no growth was taken and incubated at 37°C for 24h. With the aid of positive and negative cultures, the lowest concentration that shows no visible bacterial growth after sub-culturing was taken as MBC and MFC. Freshly prepared sterile nutrient agar was poured separately into sterile petridishes and allowed to set firmly, a loopful of mixture in the tube not showing growth was transferred to the agar in the plates and incubated for 24 hrs. At 37°C and observed for growth. The plate that did not show growth was recorded as MBC and MFC against the test organisms, respectively.

Statistical Analysis

One way analysis of variance (ANOVA) was used to statistically analyze all the data obtained from research using Statistical Package for Social Sciences (SPSS) Software version 22 and results were recorded.



(a) Dried, weighed extracts



(b) Soaking of extracts



(c) Crude extracts



(d) Drying of extract in rotatory evaporator



(e) Drying of ethanolic extracts on hot plate



(f) Drying of aqueous extracts In hot water bath



(g) Phytochemical tests



(h) Macfarland standard



(i) Column packed with silica gel



(j) Elution of fractions



(k) Frictions elluteded from column

Figure 2: Extraction Processes of *Ocimum gratissimum* leaves

Results

The results of the analysis carried out on the *Ocimum gratissimum* leaves extract are presented as follows:

NAME OF PLANT	SOLVENT USED FOR EXTRACTION	WEIGHT OF POWDERED EXTRACT	WEIGHT OF CRUDE EXTRACTS
<i>Ocimum gratissimum</i>	Water	300g	147.0g
<i>Ocimum gratissimum</i>	Ethanol	300g	130.1g

Table 1: The Yield of Extracts of *Ocimum gratissimum*


Extraction Solvents	<i>Ocimum gratissimum</i>	
	AQUEOUS	ETHANOL
Phytochemicals		
Tannins	+	++
Alkaloids	+	+++
Flavonoids	++	+++
Saponins	+	+++
Glycosides	-	+
Steroids	-	+
Phenolics	++	+++
carbohydrates	+	-

Key: (+++) = Highly present,
 (++) = Moderately present,
 (+) = Slightly present, (-) = Absent


Table 2: Phytochemical Composition of Aqueous and Ethanolic Extracts of *Ocimum gratissimum* Leaves

Thin Layer Chromatography (TLC)

Thin-layer chromatographic study revealed that the ethanolic and aqueous extracts of *O. gratissimum* has a maximum of 5 bands. The solvent front was 7.0cm with RF of between 0.14-0.53 as shown in figure bellow.



a. Aqueous extract



b. Ethanolic extract

Figure 3: Thin-layer chromatographic bands of *Ocimum gratissimum* leaves extract

FRACTIONS OF OCIMUM GRATISSIMUM LEAVES EXTRACT ELUTED FROM COLUMN CHROMATOGRAPHY

During column chromatography, 12 fractions were eluted from *O. gratissimum* leaves extract. For the, bioactive compounds the distance moved by the solvent front was 8.0cm with Rf value of between 0.06- 0.44 as shown in Tables 3 and Figures 4.

No. of Bands	Fraction	Rf values	Colour of bands	Spraying Reagent	Colour of bands appeared	Phytochemicals Detected
1	F1	0.31	Light green	Ammonia solution	Dark grey	Flavonoid
2	F2-3	0.43	Dark green	Conc. HCl	Dark brown	Saponins
3	F4-6	0.29	Orange	Dragendorff's reagent	Orange	Alkaloid
4	F7-9	0.14	Light yellow	Ammonia solution	Yellow	Flavonoid
5	F10-12	0.17	Light grey	FeCl ₃	Red	Phenol

Table 3: ELUTION PROFILE OF COLUMN CHROMATOGRAPHY OF ETHANOLIC LEAF EXTRACTS OF *OCIMUM GRATISSIMUM* IN ON MOBILE PHASE OF N- HEXANE: ETHYL ACETATE: WATER (4:2:1)




Figure 4: TWELVE FRACTIONS OF *O. GRATISSIMUM* LEAVES EXTRACT ELUTED FROM COLUMN CHROMATOGRAPHY

SUCEPTIBILITY OF THE DIFFERENT FRACTIONS AGAINST TEST ORGANISMS

The antimicrobial activity of the extract of *Ocimum gratissimum* were located at Rf values of between 0.06- 0.44. F1 has the highest activity on *Escherichia coli* with zone of inhibition of 20.00mm, 9.11mm on *Staphylococcus aureus* and 6.01mm against *Candida albicans*. F 2-3 shows the antimicrobial activity on *Escherichia coli* with zone of inhibition of 15.00mm, 8.01mm on *Staphylococcus aureus* and 7.00mm on *Candida albicans*, F4-6 also showed activity on *Escherichia coli* with zone of inhibition of 6.03mm, 4.02 against *S.aureus* and 2.00mm against *Candida albicans* while F 7-9 did not show activity against *Escherichia coli*, *Staphylococcus aureus* and *Candida albicans*, F10-12 also did not show any activity on any of the organisms.


Ocimum gratissimum

Fraction	<i>E. coli</i>	<i>Staph aureus</i>	<i>Candida albicans</i>
F1	20.00	9.11	6.01
F 2-3	15.00	8.01	7.00
F4-6	6.03	4.02	2.00
F 7-9	-	-	-
F10-12	-	-	-

Key:

(-) = No inhibition F = Fraction

Table 4: THE ANTIMICROBIAL ACTIVITY OF THE DIFFERENT FRACTIONS OF *O.gratissimum* LEAVE EXTRACT AGAINST THE THREE UROPATHOGENS



Key: F = Fraction

Figure 5: Inhibition zones of the fractions from column chromatography on the uropathogens (a. *E.coli*, b. *S.aureus*, c. *C. albicans*) respectively

Identification of Test Organisms

Each organism was identified using their specific biochemical tests. Confirmatory tests carried out to identify the isolates were Gram staining reaction, indole test, catalase test, coagulase test and germ tube test. It was observed that *E. coli* was negative for Gram staining reaction, positive for indole test, negative for catalase, coagulase tests, and germ tube test. *Staphylococcus aureus* was catalase positive, positive for gram staining reaction, negative for indole test, and coagulase positive and germ tube test negative. *C. albicans* was only positive for germ tube test negative for Gram staining reaction, indole test, catalase production and coagulase test as shown in table 5.

Gs	In	Cat	Cao	Gtt	Identity of isolates
-	+	-	-	-	<i>E. coli</i>
+	-	+	+	-	<i>Staph aureus</i>
+	-	-	-	+	<i>C.albicans</i>

Key: Gs-Grams staining Reaction, In-indole; Cat- catalase; Coa-coagulase, Gtt-germ tube test, +: Positive -: Negative.

Table 5: IDENTIFICATION OF UTI ISOLATES

Antimicrobial Activity of Aqueous Extract of The Leaves on The Three Uropathogens

The aqueous extract of *Ocimum gratissimum* leaves extract showed weak antimicrobial activity on the test organisms with 3.5 ± 0.5 mm as the highest zone of inhibition at a concentration of 500mg/ml and 1.4 ± 0.0 mm as the lowest zone of inhibition at a concentration of 31.25mg/ml.


Table 6 shows the susceptibility of aqueous extract of *Ocimum gratissimum* leaves extract on the three UTI isolates and also on standard strains. It was observed that all the three isolates were inhibited at concentration of 500mg/ml with varying zones of inhibitions. Also at 250mg/ml and 125mg/ml all the three isolates were also inhibited at varying zones of inhibitions but at concentration of 62.5mg/ml, only *S. aureus* was inhibited with zone of inhibition of 0.5 ± 0.7 c while the other test organisms were not inhibited. At 31.25mg/ml none of the organisms was inhibited. The LSD was calculated against each organisms with *S.aureus* having the highest LSD of 1.80 and *E.coli* ATCC11775 having the least LSD of 0.74. Means tagged with different letter alphabet are significantly different at $P=0.05$ example 5.5 ± 0.5 a and 2.1 ± 0.1 b against *E.coli* while Means tagged with the same letter of alphabet are not significant at $P=0.05$ example 2.1 ± 0.1 b and 1.6 ± 0.6 b against *E.coli*.

conc(mg/ml)	<i>E.coli</i> (mm)	<i>S.aureus</i> (mm)	<i>C.albicans</i> (mm)	<i>E.coli</i> ATCC11775 (mm)	<i>S.aureus</i> ATCC 6538 (mm)	<i>C.albicans</i> ATCC10231 (mm)
500	3.5 ± 0.5 a	3.0 ± 1.1 a	2.0 ± 0.0 a	3.9 ± 0.1 a	3.1 ± 0.1 a	2.5 ± 0.5 a
250	2.1 ± 0.1 b	2.1 ± 0.0 b	1.8 ± 0.1 b	2.0 ± 0.0 b	2.0 ± 0.0 b	1.9 ± 0.1 a
125	1.6 ± 0.6 b	1.9 ± 0.1 b	1.4 ± 0.1 c	1.4 ± 0.0 b	1.8 ± 0.1 c	1.7 ± 0.1 a
62.5	-	0.5 ± 0.7 c	-	-	-	-
31.25	-	-	-	-	-	-
LSD	1.21	0.8	0.18	0.74	0.18	0.82

Footnote: Means tagged with different letter alphabet are significant at $P=0.05$

Key: - = No inhibition

Table 6: Suceptibility Of Aqueous Extract of *Ocimum Gratissimum* Leaf on The Three Uti Isolates and Standard Strains



a) *O.gratissimum* against *E. coli* b) *O.gratissimum* against *S.aureus* c) *O.gratissimum* against *C.albicans*



d) *O. gratissimum* against *E. coli* ATCC 11775 e) *O. gratissimum* against *S. aureus* ATCC 6538 f) *O. gratissimum* against *C. albicans* ATCC 10231

Figure 6: Suceptibility of aqueous extract of *O. gratissimum* leaves on the UTI Pathogens with their corresponding standards

ETHANOLIC EXTRACTS OF *Ocimum gratissimum* LEAF

The ethanolic extract of *Ocimum gratissimum* leaves extract exhibited strong antimicrobial activity *Escherichia coli*, *Staphylococcus aureus* and *Candida albicans*. It also inhibited the organisms with maximum zones of inhibitions of 23.5 ± 1.5 , 23.6 ± 0.6 and 18.25 ± 1.3 mm, respectively at contraction of 500 mg/ml.

The susceptibility of ethanolic extract of *Ocimum gratissimum* leaves extract on the three UTI isolates and standard strains is shown in Table 7. It was observed that all the three isolates were inhibited at concentration of 500 mg/ml with varying zones of inhibitions- 23.5 ± 1.5 a, 23.6 ± 0.6 a, 18.25 ± 1.3 a, 22.0 ± 1.9 a, 24.6 ± 0.4 a and 19.0 ± 1.0 on *Escherichia coli*, ATCC11775, *Staphylococcus aureus* ATCC 6538 and *Candida albicans* ATCC10231 respectively. Also at 250 mg/ml and 125 mg/ml all the three isolates were inhibited at concentration of with varying zones of inhibitions but at concentration of 62.5 mg/ml, only *E. coli* isolate, and *E. coli* strain- ATCC11775 were inhibited with zones of inhibition of 4.1 ± 4.1 b and 3.5 ± 3.5 b respectively while the other test organisms were not inhibited. At 31.25 mg/ml, none of the organism was inhibited. The LSD were calculated against each organisms with *C. albicans* ATCC10231 having the highest LSD of 10.00 and *S. aureus*, having the least LSD of 3.30. Means tagged with different letter alphabet are significantly different at $P=0.05$ as in table 7.

Conc (mg/ml)	<i>E. coli</i> isolate (mm)	<i>S. aureus</i> isolate (mm)	<i>C. albicans</i> isolate (mm)	<i>E. coli</i> strain- ATCC11775 (mm)	<i>S. aureus</i> strain- ATCC 6538 (mm)	<i>C. albicans</i> strain- ATCC10231 (mm)
500	23.5 ± 1.5 a	23.6 ± 0.6 a	18.25 ± 1.3 a	22.0 ± 1.9 a	24.6 ± 0.4 a	19.0 ± 1.0 a
250	19.7 ± 0.4 a	19.2 ± 1.3 b	12.1 ± 2.1 b	19.3 ± 0.1 a	18.7 ± 1.7 b	7.5 ± 5.5 b
125	15.3 ± 3.3 a	8.5 ± 1.5 c	6.6 ± 0.6 c	14.6 ± 3.4 a	9.2 ± 1.3 c	3.5 ± 2.5 b
62.5	4.1 ± 4.1 b	-	-	3.5 ± 3.5 b	-	-
31.25	-	-	-	-	-	-
LSD=	8.87	3.30	4.10	8.55	3.51	10.00

Footnote: Means tagged with different alphabet are significant at $P=0.05$

Key: (-) = No inhibition

Table 7: Antimicrobial Activity of Ethanolic Extract of *Ocimum Gratissimum* Leaf on The Three Uti Isolates And Standard Strains



a) *O. gratissimum* against *E. coli* b) *O. gratissimum* against *S. aureus* c) *O. gratissimum* against *C. albicans*



d) *O. gratissimum* against *E. coli* ATCC 11775 e) *O. gratissimum* against *S. aureus* ATCC 6538 f) *O. gratissimum* against *C. albicans* ATCC 10231

Figure 7: Susceptibility test of ethanolic extract of *O. gratissimum* leaves on the Uropathogens with their corresponding standards strains

Positive and Negative Controls

Positive (Nitrofurantoin) was used on *E. coli*, Streptomycin was used on *S. aureus* while Fluconazole was used on *C. albicans*. The zones of inhibitions were 25.6 ± 0.5 , 20.0 ± 0 and 18.0 ± 1.0 on *E. coli*, *S. aureus*, and *C. albicans* respectively at the concentration of 500mg/ml. The zones of inhibitions for the standard strains were; 21.1 ± 0.1 , 20.5 ± 2.5 and 18.0 ± 1.0 at the concentration of 500mg/ml for *E. coli* ATCC 11775, *S. aureus* ATCC 6538, and *C. albicans* ATCC 10231 respectively. At the concentration of 250mg/ml and 125mg/ml all the three isolates were inhibited at concentration of with varying zones of inhibitions but at concentration of 62.5, only *E. coli* was inhibited with zones of inhibition of 1.00mm. At 31.25 none was inhibited.

Comparing the zones of inhibition of the antibiotics to that of plant extracts, it was observed that the highest zone of inhibition of the control was 25.6 ± 0.5 mm while the highest zone of inhibition of the Plant extracts was 24.7 ± 0.5 mm. There is statistical difference as the difference between the zones of inhibition of the plant extract to that of the antibiotics was just as little as 0.9 ± 0.0 mm. The negative control (sterile distilled water) did not show any antimicrobial activity in both the aqueous and ethanolic leave extracts as presented in plates below.

Conc. of positive controls	<i>E. coli</i>	<i>E. coli</i> ATCC11775	<i>S. aureus</i>	<i>S. aureus</i> ATCC 6538	<i>C. albicans</i>	<i>C. albicans</i> ATCC10231
500	25.6 ± 0.5	20.0 ± 0	18.0 ± 1.0	21.1 ± 0.1	20.5 ± 2.5	18.1 ± 0.0
250	18.6 ± 1.0	16.6 ± 1.0	10.0 ± 1.0	10.0 ± 1.0	12.0 ± 1.0	11.5 ± 0.5
125	8.5 ± 0.5	5.5 ± 0.5	4.3 ± 0.5	3.5 ± 0.5	2.2 ± 0.0	2.0 ± 0.0
62.5	1.0 ± 0.0	-	-	-	-	-
31.25	-	-	-	-	-	-
Negative Control (Sterile Distilled water)	-	-	-	-	-	-

Key: = No inhibition

Table 8: SUSCEPTIBILITY TEST ON POSITIVE AND NEGATIVE CONTROLS
Positive controls Nitrofurantoin Streptomycin Fluconazole




Figure 8: Positive and negative controls

NOTE: Positive control – Antibiotics Negative control - Distilled water

Minimum Inhibitory Concentrations (Mic) of The Extract on The Isolates

The Minimum Inhibitory Concentration (MIC) was determined using broth dilution method. The organisms were observed for their ability to produce visible growth on series of broth dilutions containing diluted antimicrobial agent. It was observed that each microorganism has a level of antimicrobial agent which inhibits their growth. The lowest concentration of antimicrobial agent in (mg/ml) that prevents the appearance of visible growth of a microorganism within 24hours was taken as the MIC.

All the microorganisms (E coli, S. aureus and C.albicans) used in this research were inhibited at varying concentrations of between 500(mg/ml) and 31.25(mg/ml). The minimum inhibitory concentrations of the ethanolic extract showed remarkable activities on the test organisms. It was equally observed that the extract has MIC of 62.5mg/ml on all the isolates except C.albicans which was inhibited at concentration of 125 mg/ml as shown in table 9.

Clinical/typed strains	500(mg/ml)	250 (mg/ml)	125(mg/ml)	62.5(mg/ml)	31.25 (mg/ml)	ESC control	MIC
S. aureus ATCC 6538	+	+	+	-	-	NG	62.5
S.aureus	+	+	+	-	-	NG	62.5
E.coli ATCC 11775	+	+	+	-	-	NG	62.5
E.coli	+	+	+	-	-	NG	62.5
C. albicans ATCC 10231	+	+	-	-	-	NG	125
C.albicans	+	+	-	-	-	NG	125

Key: + = Growth - = No growth ESC = Extract sterility control

Table 9: Mic of Ethanolic Extracts of Ocimum Gratissimum Leaves on The Isolates
Mean Diameter of zone of inhibitions shown by different concentration of the extract (mm)

Minimum Bacteriostatic Concentration (Mbc) and Minimum Fungistatic Concentration (Mfc) of The Extracts

It was observed that the extracts have MBC of 62.5mg/ml on both the isolates and their standard strains.

Clinical/typed strains	500(mg/ml)	250 (mg/ml)	125(mg/ml)	62.5(mg/ml)	31.25(mg/ml)	ESC	MIC
S.aureus strain-ATCC 6538	+	+	+	-	-	-	62.5
S.aureus isolate	+	+	+	-	-	-	62.5
E.coli strain-ATCC 11775	+	+	+	-	-	-	62.5
E.coli isolate	+	+	+	-	-	-	62.5
C.albicans strain-ATCC 10231	+	+	+	-	-	-	62.5
C.albicans isolate	+	+	+	-	-	-	62.5

Key: + = Growth - = No growth ESC = Extract sterility control

Table 10: The Mbc and Mfc Results of The Ethanolic Extract on The Organisms
Mean Diameter of zone of inhibitions shown by different concentration of the extract (mm)

Discussion

Several phytochemical compounds which exhibited antimicrobial activities were extracted from the leaves of *O.gratissimum* and this suggest that urinary tract infections can be reduced or treated with natural products (Flores-Mireles et al., 2015) such as the leaves of *O. gratissimum* which may aid in the reduction of the prevalence of UTI if well harnessed. The phytochemical tests of the aqueous and ethanolic extracts showed that the leaves contains Tannins, Alkaloids, Flavonoids, Saponins, Glycosides, Steroids, Phenolics, and carbohydrates. These compounds have been documented by other researchers in previous studies (Sofowora, 1993). The presence of these phytochemicals in *O. gratissimum* leaves extract is an indication

that the plant has pharmacological importance which may be used as drug in treating UTI pathogens (Adebolu & Salu, 2005).

This is in affirmation to the result obtained by Cowan (1999) who reported that plant extracts possess different phytochemicals. In this study it was observed that alkaloids, flavonoids, saponins and phenol were higher in concentration than the other phytochemical compounds. This is in accordance to the work of Nweze et al. (2009) who also observed the presence of alkaloids, flavonoids, saponins and phenol in higher concentration, this maybe as a result of using the same method

of extraction. Sterols was equally extracted from the leaves of *O. gratissimum* and present sterols have also been reported to have antibacterial properties in other studies (Cowan, 1999; Okwu, 2001). The presence of alkaloids and saponins in the present work is in agreement with the opinion of Kasolo et al. (2010) who noted that saponins and alkaloids were two active constituents in plants.

It was equally documented by other studies that alkaloids are nitrogenous compounds occurring naturally which are commonly found to have antimicrobial properties like in this study due to their ability to intercalate with DNA of microorganisms. However, the presence of saponin in this plant is in agreement with the previous findings of Yusha'a et al. (2011) who also extracted saponin in his study (Yusha'a et al., 2011). The activity of plant extracts against bacteria have been studied for years, but in more intensified way during the last three decades. During this period, numerous antimicrobial screening have been published based on different traditional regions like; Chinese, African and Asian countries who have used some of these plant as drugs (Suffredim et al., 2004).

The aqueous extracts of *O. gratissimum* leave extracts showed weak antimicrobial activities and this maybe as a result of the extraction solvents used, knowing fully well that water is more polar than other compounds, so non polar compounds may not have been extracted from the leaves. This means that a lot of bio-active compound may not have been extracted by water. This result is contrary to the result of Lapornik et al. (2005) who reported that water is a better extraction solvent as compared to ethanol because in this study, water was not the best solvent for extraction. However, it is in accordance with the findings of Das et al. (2010) who reported that water soluble flavonoids have no antimicrobial significance and water soluble phenolics are only important as antioxidant compounds. Though traditional healers use primarily water to extract plant but organic solvents have been found to give more consistent antimicrobial activity compared to water extract (Lapornik et al., 2005).

In this present study, the results of the antimicrobial properties of the plant extracts on different organisms varied depending on the organism tested and the concentration of the extracts, it was observed that the zone of inhibition decreased with decrease in concentration of the extracts and increase with increased in concentration of the extracts. This is comparable to previous studies (Taylor et al., 2001; Farnsworth, 1993) which recorded that active compound may be present in insufficient quantities in extracts which may or may not show antimicrobial activity on pathogenic organisms. It is observed that the higher the concentration, the higher the antimicrobial activity, it is therefore suggested that in order to rule out the inability of plants extract to show antimicrobial activity, high concentration should be used (Farnsworth, 1993; Jager et al., 1993).

The ethanolic extracts exhibited stronger activity on all the UTI pathogens at varying concentrations. This may be attributed to the extraction solvent (ethanol) used or maybe the compounds

in the leaves were less polar since ethanol is polar. It should be noted that ethanol was not inhibiting the uropathogens rather it was the bioactive compounds in the ethanolic extracts doing the inhibition because all the ethanol used in the extraction was evaporated via rotary evaporator. Ethanol was considered to be more efficient in cell walls and seeds degradation which have unipolar character and cause polyphenols to be released from cells (Cowan, 1999).

Conclusion

Having done thorough investigation beyond any reasonable doubt, the leaves extract of *O. gratissimum* have exerted strong, good and promising antimicrobial activities on Uropathogens. The antimicrobial activities it shows on both the gram positive and gram negative bacteria as well as on fungi shows that it has a broad spectrum effect which is very impressive. The differences observed in the antimicrobial activities between aqueous and ethanolic extracts shows that solvent used for extraction is very paramount.

This present demonstrated that herbal medicines can also be effective as modern medicine to combat some pathogenic microorganisms if well harnessed. The actual bioactive phytochemicals in the *Ocimum gratissimum* leaves chromatographically identified are flavonoids, saponins and phenols.

It is recommendable that further study should be carried on *Ocimum gratissimum* leaves extract in order to convert the essential phytochemicals in the leaves to complimentary medicine in the treatment of urinary tract infections leading to the formulation of natural UTI drugs.

Acknowledgments

The Authors acknowledge Medical Laboratory Scientists who are contributing in alternative medicine discovery research.

Conflict of Interest

None

Financial Support

None

Ethics Statement

None.

References

1. World Health Organization. (2013). *WHO Traditional Medicine Strategy: 2014- 2023*. Retrieved from WHO traditional medicine strategy: 2014-2023
2. Ezekwesili-Ofili, J. O., Okaka, A. N. C., (2019). Herbal Medicines in African Traditional Medicine Chapter 10, IntechOpen. <http://dx.doi.org/10.5772/intechopen.80348>
3. Obeta M. U., Ikeagwulonu, R. C., Ohanube G. A. K., & Jwanse, I. R. (2020). Some Igbo Indigenous Plants with Anti-COVID-19 Properties, a Chapter. In *Alternative Medicine Update*. IntechOpen DOI: <http://dx.doi.org/10.5772/intechopen.94244>
4. Obeta, M. U., Ohanube, G. A. K., & Akram, M. (2021). The Truth about Vitamin C Enriched Plants and Vegetables used in Nigeria for the Management of COVID-19.

- Edelweiss Food Sci Tech*, 7-9.
<https://doi.org/10.33805/2765-8821.109>
5. Agbalaka, P., Matthew, G., Obeta, U., Sabulu, J., Joshua-Ojokpe, R., & Pada, N. (2021). Comparative Insecticidal Effects of Dry *Ocimum Gratissimum* (Scent Leaves) and Rambo™ Paper on Mosquitoes in Jos, Nigeria. *Bioremediation Science and Technology Research*, 9(2), 13-16. <https://doi.org/10.54987/bstr.v9i2.620>
 6. Anzaku, F. A., Obeta, K. O., Mairiga, J. P., Obeta, M. U., Ejinkaka, O. R., Akram, M., Ekpere-Ezeugwu, M. O., & Elbossaty, W. (2021). Evaluation of the Insecticidal Effects of *Ocimum Sanctum* on Mosquito. *International Journal of Pharmaceutical and Phytopharmacological Research*, 11(6), 11-17 <https://doi.org/10.51847/oRXfaNnQNG>
 7. Bhavani, T., Mohan, R. R., Mounica, C., Nyamisha, J., Krishna, A. G., Raja, R. R., & Harinadha, B. K. (2019). Phytochemical screening & antimicrobial activity of *Ocimum gratissimum* review. *J Pharmacogn Phytochem*, 8(2), 76-9. Retrieved from 8-1-533-370.pdf (phytojournal.com)
 8. Lunyera, J. D., Wang, V., & Maro, V. (2016). Traditional medicine practices among community members with diabetes mellitus in Northern Tanzania: An ethnomedical survey. *BMC Complementary and Alternative Medicine*, 16(1), 282. DOI: 10.1186/s12906-016-1262-2
 9. Nwinyi, Obinna. C., Chinedu. Nwodo, S., Ajani, Olayinka, O., Ikpo. Chinwe, O., & Ogunniran, Kehinde. O. (2009). Antibacterial effects of extracts of *Ocimum gratissimum* and *Piper guineense* on *Escherichia coli* and *Staphylococcus aureus*. *African Journal of Food Science*, 3(3), 077-081. Retrieved from Microsoft Word - Nwinyi et al pdf.doc (academicjournals.org)
 10. Dhanalakshmi, J., S. Selvi. (2013). Antibacterial Activity of Medicinal Plants used against UTI (Urinary Tract Infection) causing Pathogens. *International Journal of Research in Sciences*. pp. 01-07
 11. Chijioko, A., Nsofor, Chikezie, U.N, Azuwuiké, C.O. (2014) Antibacterial activity of *Ocimumgratissimum* leaves extract on *Escherichia coli* O157. *Pjpalgo journal of medicine and medical sciences*, Page 15-18.
 12. Bazzaz, B. S. F., Fork, S. D., Ahmadi, R., & Khameneh, B. (2021). Deep insights into urinary tract infections and effective natural remedies. *Afr J Urol*, 27(6). <https://doi.org/10.1186/s12301-020-00111-z>
 13. Nweze, E.I., & Eze, E.E. (2009). Justification for the use of *Ocimum gratissimum* L in herbal medicine and its interaction with disc antibiotics. *BMC Complementary and Alternative Medicine*, 9(37). DOI:10.1186/1472-6882-9-37
 14. Iroanya, O.O., Okpuzor, J.E., Mbagwu, H., & Ojobo, P.D. (2009). Analgesic properties of an indigenous polyherbal preparation. *The FASEB Journal*, 23(S1), 712.2-712.2. https://doi.org/10.1096/fasebj.23.1_supplement.712.2
 15. Nguéfacq, J., Dongmo, J.B.L., Dakole, C.D., Leth, V., Vismar, H.F., Torp, J., Guemdjom, E.F.N., Mbeffo, M., Tamgue, O., Fotio, D., Zollo, P.H.A., & Nkengfacq, A.E. (2009). Food preservative potential of essential oils and fractions from *Cymbopogon citratus*, *Ocimum gratissimum* and *Thymus vulgaris* against mycotoxigenic fungi. *International Journal of Food Microbiology*, 131(2-3), 151-156. <https://doi.org/10.1016/j.ijfoodmicro.2009.02.009>
 16. Sofowora, A.E. (1993). Medicinal Plants and Traditional Medicine in Africa. Spectrum Books Ltd. Ibadan, Nigeria, pp. 191-289. Retrieved from Sofowora, A. (1993) Medicinal Plants and Traditional Medicine in Africa. Spectrum Books Ltd., Ibadan, 191-289. - References - *Scientific Research Publishing* (scirp.org)
 17. Owoyele, B. V., Funsho, M. A., & Soladoye, A.O. (2005). Effect of aqueous leaves extract of *ocimum gratissimum* (sweet basil) on alloxan induced diabetic rats. *PHCOG MAG*, 1(2), 62-64. Retrieved from Effect of aqueous leaves extract of *ocimum gratissimum* (sweet basil) on alloxan induced diabetic rats. | Semantic Scholar
 18. Egesie, U. G., Adelaiye, A. B., Ibu, J. O., & Egesie, O. J. (2006). Safety and hypoglycaemic properties of aqueous leaf extract of *Ocimum gratissimum* in streptozotocin induced diabetic rats. *Niger Journal of Physiol Sci*, 21 (1-2), 31 35. DOI: 10.4314/njps.v21i1-2.53971
 19. Ayinla, M. T., Dada, S. O., Shittu, S. T., Olayaki, L. A., Akiode, A. O., & Ojulari, S. L. (2011). Anti hyperlipidemic effect of aqueous leaf extract of *Ocimum gratissimum* in alloxan induced diabetic rats. *Int Journal of Med Sci*, 3(12), 360 363. <https://doi.org/10.5897/IJMMS.9000022>
 20. Shittu, S. T., Oyeyemi, W. A., Lasisi, T. J., Shittu, S. A., Lawal, T. T., & Olujobi, S. T. (2016). Aqueous leaf extract of *Ocimum gratissimum* improves hematological parameters in alloxan-induced diabetic rats via its antioxidant properties. *International Journal of App Basic Med Res*, 6(2), 96-100. DOI:10.4103/2229-516X.179016
 21. Silva, R. M., Paredes, J. A., Moura, G. R., Manadas, B., Lima-Costa, T., Rocha, R., Miranda, I., Gomes, A. C., Koerkamp, M. J. G., Perrot, M., Holstege, F. C. P., Boucherie, H., & Santos, M. A. S. (2007). Critical roles for a genetic code alteration in the evolution of the genus *Candida*. *EMBO Journal*, 26(21), 4555-4565. <https://doi.org/10.1038/sj.emboj.7601876> (21): 4555–65.
 22. Lexa, G. Matasyoh., Josphat, C. Matasyoh., Francis, N. Wachira., Miriam, G. Kinyua., Anne, W. Thairu., M., & Titus, K. M. (2006). Chemical composition and antimicrobial activity of the essential oil of *Ocimum gratissimum* L. growing in Eastern Kenya. *African Journal of Biotechnology*, 6(6), 760-765. Retrieved from Chemical composition and antimicrobial activity of the essential oil of *Ocimum gratissimum* L. growing in Eastern Kenya | Semantic Scholar
 23. Matasyoh, L.G., Matasyoh, J.C., Wachira, F.N., Kinyua, M.G., Thairu, Muigai. A.W., & Mukiyama, T. K. (2007). Chemical Composition and antimicrobial activity of the essential oil of *Ocimum gratissimum* L. growing in Eastern Kenya. *African Journal of Biotechnology*, 6(6), 760-765. Retrieved from Matasyoh, Lexa G. 1*, Josphat C. Matasyoh2, Francis N. Wachira3, Miriam G. Kinyua4, Anne W. Thairu Muigai1 and Titus K. Mukiyama 5(2007) Chemical composition and antimicrobial activity of the essential oil of *Ocimum gratissimum* L. growing in

- Eastern Kenya. | PROF. MUKIAMA TITUS K. (uonbi.ac.ke)
24. Nakaruma, C.V., T.U. Nakaruma, E. Bando., A.F.N. Melo, D.A.G. Cortez.,B.P, DiazFilho. (1999). Antibacterial activity of *Ocimum gratissimum* L.essential Oil. Mem. Inst. OswaldoCruz, 94, 675-578.
 25. Lemos, J. A., Passos, X. S., Fernandes, O. F., Paul, J. R., Ferri, J. P., Ferri, P. H., Souza, L. K., Lemos, A. A., & Silva, K. R. M. (2005). Antifungal activity of *Ocimum gratissimum* towards *Cryptococcus neoformans*. *Journal of Ethnopharmacology*, 20, 4–13.
 26. Kolkate, C. K., A. P, Purohit, Kokhale, S. B. (2003). Qualitative chemical examination.In Text book of pharmacognosy. *Nitani Publication Pune*.Pp 108-109.
 27. Harwood, L. M, Moody C. J. (1989) Experimental Organic Chemistry: Principles and Practice Illustrated Ed. Wiley-Blackwell, Hoboken, 122-125. Retrieved from Harwood, L.M. and Moody, C.J. (1989) Experimental Organic Chemistry Principles and Practice (Illustrated Edition). Wiley-Blackwell, Hoboken, 122-125. - References - Scientific Research Publishing (scirp.org)
 28. Martin, D. F. (2013). Chromatographic Separations with Selected Supported Chelating Agents. In Column Chromatography Eds. Martin DF, Martin BB. Intech Open <https://doi.org/10.5772/55521>
 29. CLSI, Methods for Determining Bactericidal Activity of Antimicrobial Agents. Approved Guideline, CLSI document M26-A. *Clinical and Laboratory Standards Institute*, 950 West Valley Road Suite 2500, Wayne, Pennsylvania 19087, USA, 1998.
 30. Flores-Mireles, A. L., Walker, J. N., Caparon, M., & Hultgren, S. J. (2015). Urinary tract infections: epidemiology, mechanisms of infection and treatment options. *Nature reviews Microbiology*, 13 (5), 269–84. DOI: 10.1038/nrmicro3432
 31. Adebolu, T. T., & Salu, A. O. (2005). Antimicrobial activity of leaf extract of *Ocimum gratissimum* on selected diarrhea causing bacteria in South Western Nigeria. *African Journal of Biotech*, 4(7), 682-684. <https://doi.org/10.5897/AJB2005.000-3126>.
 32. Cowan, M. M. (1999). Plant products as antimicrobial agents. *Clinical microbiology reviews*, 12(4), 564 582. DOI: 10.1128/CMR.12.4.564
 33. Okwu, D. E. (2001). Evaluation of the chemical composition of indigenous spices and flavouring agents. *Global Journal of Pure Appl. Sci*, 7(3), 455-459. Retrieved from D. E. Okwu, "Evaluation of the Chemical Composition of Indigenous Spices and Flavouring Agents," *Global Journal of Pure and Applied Sciences*, Vol. 7, No. 3, 2001, pp. 455-459. - References - Scientific Research Publishing (scirp.org)
 34. Kasolo, J. N., Bimenya, G. S., Ojok, L., Ochieng, J., &Ogwal-Okeng, J. W. (2010). hemicalsandusesof Moringaoleiferaleavesin Ugandanrural communities. *Journal of Medicinal Plants Research*, 4(9), 753-757. Retrieved from Kasolo, J.N., Bimenya, G.S., Ojok, L., Ochieng, J. and Ogwal-Okeng, J.W. (2010) Phytochemicals and Uses of Moringa oleifera Leaves in Ugandan Rural Communities. *Journal of Medicinal Plants Research*, 4, 753-757. - References - *Scientific Research Publishing* (scirp.org)
 35. Yusha'a, M., Taura, D. W., Bello, B., & Abdullahi, N. (2011). *International Research of Pharmacy and Pharmacology*, 1(9), 237-241. Retrieved from Screening of *Annona squamosa* extracts for antibacterial activity against some respiratory tract isolates (scholarscentral.com)
 36. Suffredini, I. B., Sader, H. S., Gonçalves, A.G., Reis, A.O., Gales, A. C., Varella, A. D., & Younes, R. N. (2004). Screening of antibacterial extracts from plants native to Brazillian amazon Rain Forest and Atlantic Forest. *Brazillian J. Med. Biol. Res*, 37(3), 379-384. DOI: 10.1590/s0100-879x2004000300015
 37. Lapornik, B., Prosek, M., & Wondra, A. G. (2005). Comparison of extracts prepared from plant by-products using different solvents and extraction time. *Journal of Food Engineering*, 71(2) 214–222. <https://doi.org/10.1016/j.jfoodeng.2004.10.036>
 38. Das, K., Tiwari, R. K. S., & Shrivastava, D. K. (2010). Techniques for evaluation of medicinal plant products as antimicrobial agent: Current methods and future trends. *Journal of Medicinal Plants Research*, 4(2), 104-111. Retrieved from Das, K., Tiwari, R.K.S. and. Shrivastava, D.K. (2010) Techniques for evaluation of medicinal plant products as antimicrobial agent Current methods and future trends. *Journal of Medicinal Plants Research*, 4, 104-111. - References - Scientific Research Publishing (scirp.org)
 39. Taylor, J. L. S., Rabe, T., McGraw, L. J., Jager, A., & Van, Staden, J. (2001). Towards the scientific validation of traditional medicinal plants. *Plant Growth Regul*, 34(1), 23-37. Retrieved from Towards the scientific validation of traditional medicinal plants | SpringerLink
 40. Farnsworth, N. R.(1993).Biological approaches to the screening andevaluationofnaturalproducts.In:Rasoanaivo P, Ratsimamanga Urver S (Eds). *Biological evaluation of plants with reference to the Malagasy Flora* (pp. 35-43). Madagascar.
 41. Jager, A. K., Hutchings, A., Raidoo, D. M., Van, Staden, J. (1999) Screening of plants used by southern African traditional healers in the treatment of dysmenorrhoea for prostaglandin-synthesis inhibitors and uterine relaxing activity. *J. Ethnopharmacol.* 64(1), 9-14. DOI: 10.1016/s0378-8741(98)00097-x
 42. Okigbo, RN.,& Mmeka, EC.(2008)Antibacterial effects of three tropical plant extractson *Staphyl ococcus aureus*, *Escherichiacoli* and *Candidaalbicans*. *Africa Journal Trend*, 5(3), 226-229. DOI: 10.4314/ajtcam.v5i3.31277
 43. Obi, V. I., & Onuoha, C. I. (2000). Extraction and Characterization Methods of Plants and Plant Products. In: Ogbulie, J.N. & Ojiako, O. A., (Eds.), *Biological and Agricultural Techniques* (271-286). Websmedia Publishers. Retrieved from Obi, V.I. and Onuoha, C.I. (2000) Extraction and Characterization Methods of Plants and Plant Products. In Ogbulie, J.N. and Ojiako, O.A., Eds., *Biological and Agricultural Techniques*, *Websmedia Publishers*, Owerri, 271-286. - References - Scientific Research Publishing (scirp.org)

-
44. Wang, C. H., Fang, C. C., Chen, N. C., Liu, S. S., Yu, P. H., Wu, T. Y., Chen, W. T., Lee, C. C., & Chen, S. C. (2012). Cranberry-containing products for prevention of urinary tract infections in susceptible populations: a systematic review and meta-analysis of randomized controlled trials. *Archives of Internal Medicine*, 172 (13), 988–996. DOI: 10.1001/archinternmed.2012.3004
 45. Mbata, T., & Saikia, A. (2007). Antibacterial activity and phytochemical screening of crude ethanolic extract of leave of *Ocimum gratissimum* L on *Listeria monocytogenes*. *The internet Journal of microbiology*, 4(2). DOI:10.5580/1b50
 46. Noumedem, J., Mihasan, M., Lacmata, S., Stefan, M., Kuate, J., & Kuete, V. (2013) Antibacterial activities of the methanol extracts of Cameroon vegetables against Gram-negative multidrug-resistant bacteria. *BMC Complement Altern Me*, 13-26. <https://doi.org/10.1186/1472-6882-13-26>
 47. Singleton, D. R., Masuoka, J., Hazen, K. C. (2005). Surface hydrophobicity changes of two *Candida albicans* serotype B mn4 delta mutants. *Eukaryot Cell*, 4(4), 639-48. DOI: 10.1128/EC.4.4.639-648.2005
 48. Bele, A.A & Knale, A. (2015). An overview on thin layer chromatography. *International Journal of pharmaceutical sciences and research*, 2(6), 256-26.
 49. Ijeh, I. I., Omodamiro, O. D., & Nwanna, I. J. (2005) Antimicrobial effects of aqueous and ethanolic fractions of two spices, *Ocimum gratissimum* and *Xylopias aethiopica*. *African J Biotech*, 4(9), 953-956. <https://doi.org/10.5897/AJB2005.000-3193>.

Copyright: ©2022. Uchejeso Mark Obeta. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.