

Impotence and Lack of Sexual Desire, Their Relation to Patients With Diabetes

Ibrahim Abdelrahim Ibrahim Humaida

Associate Professor of Mental Health. Faculty of Arts / Department of Psychology, Omdurman Islamic University- Sudan

*Correspondence author

Ibrahim Abdelrahim Ibrahim Humaida

Associate Professor of Mental Health.
Faculty of Arts / Department of Psychology
Omdurman Islamic University
Sudan

Submitted : 8 Nov 2022 ; Published : 29 Nov 2022

Citation: Ibrahim Abdelrahim Ibrahim Humaida(2022). Impotence and Lack of Sexual Desire, Their Relation to Patients With Diabetes. J Psychol Neurosci; 2022; 4(4):1-6.

Abstract

The main objective of this study was to investigate the prevalence of sexual disorders in diabetics, owing to diabetes constitutes a growing public health problem, leading to a variety of dysfunctions such as cardiovascular, psychological, and sexual dysfunctions, that is why Diabetes is a well-known cause of sexual disorders, with prevalence rates approaching 50% in both type 1 and type 2 diabetes, but the determinants of sex dysfunction in diabetic men as a result of the principal cardiovascular risk factors, such as hypertension, and also overweight obesity, in addition to metabolic syndrome, smoking, and sedentary lifestyles. Moreover, sexual disorders considered as important predictors of the development of major complaints in diabetic patients, on the other hand, the debate as to whether diabetes link with sexual dysfunction or not, is an issue of controversy, moreover, diabetic women suffer from the same neurovascular complications that contribute to the pathogenesis of SD in men, however, results of sexual functioning of diabetic women are less conclusive. Conclusion: extending beyond the specific effects on sexual dysfunction in men and women with diabetes, the adoption of these measures promotes a healthier life and increased well-being, which in turn, may help to reduce the burden of sexual dysfunction.

Keywords : Diabetics, Sexual Dysfunction, Impotence, Desire

Introduction

The causes of erectile dysfunction in men with diabetes are complex and involve impairments in nerve, blood vessel, and muscle function (Tamás & Kempler, 2014).

To get an erection, men need healthy blood vessels, nerves, male hormones, and a desire to be sexually stimulated. Diabetes can damage the blood vessels and nerves that control erection. Therefore, and more likely, diabetics might have normal amounts of male hormones, have the desire to have sex, but still may not be able to achieve a firm erection (International Diabetes Federation [IDF], 2015).

Diabetes has been associated with sexual dysfunction both in men and in women. Diabetes is an established risk factor for sexual dysfunction in men; a threefold increased risk of erectile dysfunction (ED) found in diabetic compared with non-diabetic men. Among women, the evidence regarding association between diabetes and sexual dysfunction is less conclusive, although most studies have reported a higher prevalence of female sexual dysfunction (FSD) in diabetic women as compared with non-diabetic women (Campos, 2012).

Diabetic men showed more probability of having an impotency than men without diabetes; moreover, the age- adjusted risk of

sexual impotence found in diabetic men compared with those without diabetes. In addition, it is estimated that the worldwide prevalence of impotence will rise to 322 million cases by the year 2025. Several cross-sectional and longitudinal studies showed an association between ED and most of the cardiovascular risk factors, such as diabetes, smoking, hypertension, metabolic syndrome, as well as depression, lower urinary tract symptoms, and poor health state. Moreover, ED is a the marker of significantly increased risk of CVD, CHD, stroke, and all-cause mortality (Rahman et al., 2007).

The Scope and Statement of the Problem

Epidemiological studies suggest that both type 1 and type 2 diabetes are associated with an increased risk of ED, which is reported to occur in $\geq 50\%$ of men with diabetes worldwide. In the Massachusetts Male Aging Study, diabetic men showed a threefold probability of having ED when compared to men without diabetes; moreover, the age-adjusted risk of ED doubled in diabetic men when compared to those without diabetes (Fox et al., 2007). Most of the studies that described the prevalence of ED in diabetes did not distinguish between type 1 and type 2 diabetics.

Two studies reported a similar likelihood of having ED among both type 1 and type 2 diabetic men, whereas another report

showed a higher risk of developing ED in men with type 1 diabetes. The occurrence of ED is 10 – 15 years earlier in men with diabetes; moreover, ED is more severe and less responsive to oral drugs in diabetes, leading to reduced quality of life (Preis et al., 2009).

Aim

To explore the effects of diabetes on sexuality, evaluate the impact of diabetes on sexual function, and assess the conventional and novel treatment approaches based on recent studies.

Literature Review

Advanced age and longer duration of diabetes have been associated with an increased risk of ED in diabetic patients (Lu et al., 2009). Whether hyperglycemia is a risk factor for the development of ED in diabetic men is still not clear (Enzlin et al., 2009). Some observational studies have shown an association between poor glycemic control, expressed by elevated levels of glycated hemoglobin (HbA1c), and ED, whereas other studies did not report any association (Maseroli et al., 2016). The different methodological approaches used in the different studies may explain, at least in part, these divergent results (Phillips & Phillips, 2015). Moreover, diabetes is commonly associated with hypertension, hyperlipidemia, overweight and obesity, metabolic syndrome, smoking, sedentary lifestyles, and autonomic neuropathy, which are recognized as risk factors for ED. Both microvascular and macrovascular diabetic complications also increase the risk of ED in diabetic men (Olarinoye & Olarinoye, 2008). The use of several medications frequently assumed by diabetic patients, such use of antihypertensive drugs (β - blockers, thiazide diuretics, and spironolactone), psychotropic drugs (antidepressants), and certain fibrates, have all been associated with an additive deleterious effect on diabetic ED (Giugliano et al., 2010). A moderate consumption of alcohol (not more than 5% of the total daily caloric intake, or ≤ 7 alcoholic drinks per week) may exert a protective effect on ED in both the general population and in diabetic men (Johannes et al., 2017).

The Underlying Causes of Impotence in Diabetics

The pathogenesis of impotence in diabetes is due to many factors, as it depends on both psychological and organic factors, as well as psychological and relationship issues, which often coexist (Hackett et al., 2016). Heart disease, neuropathy, visceral adiposity, insulin resistance, and disturbed gonads represent the proposed mechanisms of erectile disorder in diabetic patients (Esposito et al., 2010).

Erectile dysfunction ED, which is an inability to get or maintain an erection strong enough or sex, is common in men who are suffering from diabetes, especially those with type 2. It can also stem from damage to nerves and blood vessels caused by poor long-term blood sugar control (Seftel et al., 2004). As mentioned, several cardiovascular risk factors associated with diabetes contribute to the genesis of penile arterial insufficiency: all of them converge on endothelial dysfunction, which represents the common denominator leading to vascular ED (Kizilay et al., 2017).

The chronic insult of hyperglycemia on the endothelium results in endothelial dysfunction, which is suggested as the link between ED and CVD (Lewis et al., 2010). A diagnosis of ED regarded as a sentinel event that should prompt the investigation of coronary artery disease (CAD) in asymptomatic diabetic men. Endothelial dysfunction in diabetes is manifested as the decreased bioavailability of nitric oxide (NO), resulting in insufficient relaxation of the vascular smooth muscle (Lewis et al., 2010). The potential mechanisms involved in endothelial dysfunction include the accumulation of advanced glycation end products; increased levels of oxygen free radicals that reduce the bioavailability of NO; impaired endothelial and neuronal NO synthesis, expression, and activity; and an imbalance between the vasoconstrictive and vasorelaxation intracellular pathways favoring increased vasoconstriction (Miner et al., 2012).

Microvascular disease determines ischemic damage in the distal circulation and autonomic and peripheral neuropathy (Esposito et al., 2009). Both somatic and autonomic neuropathies may contribute to diabetes-induced erection due to the impairment of sensory impulses from the penis to the reflexogenic erectile center and reduced or absent parasympathetic activity necessary for relaxation of the smooth muscle of the penis (Esposito et al., 2005 ; Isidori et al., 2017).

Insulin resistance and visceral adiposity, which are both distinctive clinical traits of type 2 diabetes, are associated with inflammatory state that results in the decreased availability and activity of hormone, leading to ED in overweight and obese diabetic men (Bellastella et al., 2013 ; Meeking et al., 2016 ; Ponholzer et al., 2005).

Subnormal testosterone concentrations found in 25% of men with type 2 diabetes in association with inappropriately low luteinizing hormone (LH) and follicle-stimulating hormone (FSH) concentrations (Go et al., 2013). Testosterone regulates nearly every component of erectile function, from pelvic ganglions to smooth muscle, and to the endothelial cells of the gonads (Giuliano et al., 2004). It also modulates the timing of the erectile process, which occurs as a function of sexual desire, coordinating penile erection with sex (Maria et al., 2014). It is still unclear what level of testosterone needed for good erectile function; however, evidence derived from clinical and molecular studies supports the use of testosterone replacement in patients with erectile disorder, although the benefit–risk ratio is uncertain in advanced age (Demir et al., 2009 ; Turek et al., 2013 ; Dong et al., 2011).

The Prevalence of Sexual Disorders in diabetic women

Sexual disorders reported in women with diabetes include the reduction or loss of sexual interest or desire, arousal or lubrication difficulties, and loss of the ability to reach orgasm (Araujo et al., 2009 ; Thorve et al., 2011).

FSD has been associated with both type 1 and type 2 diabetes. A recent meta-analysis that included 26 studies, 3,168 diabetic women, and 2,823 controls showed that FSD is more frequent, and is associated with a lower Female Sexual Function Index

(FSFI) score in diabetic women than in controls (Corona et al., 2013). In particular, the risk for FSD was 2.27 (95% confidence interval [CI]: 1.23–4.16) and 2.49 (95% CI: 1.55–3.99) in type 1 and type 2 diabetic women, respectively (Chew et al., 2014). Having a high sugar levels for a long period of time can cause sexual problems, women with diabetes are more at risk of having sex impairment. Diabetes can damage blood vessels and nerves that supply sex organs. This can restrict the amount of blood flowing to sexual organs, so the patient can lose sensation to sexual stimuli. This means she has a difficulty getting aroused, vaginal dryness, and painful sex as well. Furthermore, damage occurs throughout the body can affect the ability to experience sexual stimulation, arousal, and the release of vaginal lubricant (Al-Hunayan et al., 2007). Interestingly, an increased risk of having sexual intercourse found in premenopausal women with diabetes, owing to urinary tract infections, which can overall impact sexual performance (Rosen et al., 2009). Moreover, at meta-regression, among the independent variables, only BMI was significantly associated with the FSFI effect size ($P=0.005$), suggesting that the higher frequency of FSD and lower FSFI score found in diabetic women may be related to body weight (Corona et al., 2015). Several studies have already shown an increased prevalence of FSD in women affected by obesity and metabolic syndrome. Studies that have focused on type 1 diabetic women have provided a valid opportunity to investigate the role of diabetes on sexual function, independent of other associated defects (Chew et al., 2013). In type 1 diabetic women, FSD appears to be correlated mainly to psychological factors, such as depression, anxiety, and marital status. Results from a large prospective study of 625 women with type 1 diabetes showed that depression was the major predictor of sexual dysfunction (Heruti et al., 2007).

Studies examining FSD in individuals with type 2 diabetes are less conclusive and limited by small study sizes; the determinants of sexual function in type 2 diabetes include age, duration of diabetes, menopause, vascular complications, and psychological complaints. In one large study that evaluated 613 diabetic women and 524 controls, found that the longer duration of diabetes, older age, higher BMI, the presence of CVD, and the presence of diabetic complications was significantly associated with worse sexual function. In a study by Esposito et al, found that metabolic syndrome an independent predictor of FSD in 595 type 2 diabetic women, although only depression and marital status were the strongest independent factors associated with FSD (Chew et al., 2008).

Diabetes-induced vascular and nerve dysfunctions may impair the sexual response by producing structural and functional changes in the female genitalia. Studies in animals showed that diabetes affect arousal and orgasmic sexual responses by inducing impaired relaxation responses of the vaginal tissue to almost all transmitter systems, decreasing nerve-stimulated clitoral and vaginal blood flow, producing diffuse fibrosis of the clitoris and vaginal tissues, and reducing the muscular layer and epithelial thickness in the vagina (Foresta et al., 2009). Vascular abnormalities, including atherosclerotic damage and diabetes-induced endothelial dysfunction, may be

responsible for reducing the engorgement of the clitoris and for reducing lubrication of the vagina, leading to decreased arousal and dyspareunia during sexual intercourse. Diabetic neuropathy may further contribute to the pathogenesis of sexual dysfunctions by altering both the normal transduction of sexual stimuli and the triggered sexual response (Hackett et al., 2016).

It has been hypothesized that FSD may be the consequence of an imbalance in the hormonal levels of diabetic women, as indicated by epidemiological studies showing a correlation between alterations in the levels of androgens, estrogens, as well as sex hormone-binding globulin and sexual problems in diabetic women. Moreover, several endocrine impairments that may be associated with diabetes, such as thyroid disorders, hypothalamic–pituitary dysfunctions, and polycystic ovarian syndrome, may further contribute to sexual dysfunctions in these women (Bal et al., 2015).

In conclusion, psychological concerns may play a significant role in the development of FSD in both type 1 and type 2 diabetes. This is in line with the complex nature of female sexuality, which is largely dependent on psychological and cultural factors, even more so than male sexuality (Thorve et al., 2011).

Methodology

In this analytical study, the researcher adopted mixed approaches in data collection that concern spreading of sexual problems among the diabetics. Quantitative data collection instruments including obtaining data from available sources, such as electronic health records, systematic reviews, clinical data, and clinical case reports. Qualitative data, which is a method that a researcher can record the behaviors of his/her research respondents directly in real life settings (naturalistic observation) at their own physical environments. Besides, narratives.

Results

1. Male and female sexual dysfunctions are a significant complication of diabetes.
2. Men were more likely to express a lack of interest in sex if they had diabetes.
3. Men also were more likely to suffer erectile dysfunction if they had diabetes.
4. Women as well as men with diabetes reported a higher rate of orgasm difficulty, including
5. climaxing too quickly for men, or not at all, which was reported by both men and women.

Discussion

Deterioration in sexual functioning is one of the major and serious complications of diabetes. This common metabolic disorder not only affects sexuality through cardiovascular and nerve damage but also has psychological aspects. In men, the primary complications are erectile dysfunction, ejaculatory dysfunction, and loss of libido. Women similarly experience sexual problems, including decreased libido and decrease in arousal and lubrication resulting in painful intercourse, and

loss of orgasm.

As to spreading of sexual problems among men and women suffering from diabetes, women are characterized by lack of libido or low sexual desire, owing to uncontrolled level of sugar in their bodies. Also, lack of orgasm due to neuropathy.

Diabetes has negative effects on both general physical health, and sexuality in particular among patients. It is noticeable that men with diabetes type 2, are reported to have retarded ejaculation, as a result of disease complication. Despite the fact that gender difference found in sexual disturbance related to diabetes, there is common clinical feature among patients all over the world.

Conclusion

Diabetes mellitus (DM) is a systemic disease that is considered to play a principal role in the etiology of sexual dysfunction. The impact of neurologic, psychogenic and vascular factors, usually combined in the pathogenesis of related complications were demonstrated in a large number of studies. Numerous medical researchers have shown that, prevalence of sexual dysfunction in men because of the degenerative effects of diabetes on vascular and neurologic structures ranges from 35 to 75%. It also documented that the etiology of ED may be neurologic in up to 82% of diabetic men. However, sexual dysfunction in diabetic women at neurological factors has not investigated in details, because female sexual dysfunction is still unexplained as male sexual dysfunction. Although cranial neuropathy and cardiovascular autonomic neuropathy may be implicated in sexual performance in both males and females.

There are some other disorders as co- morbidity that can lead to appearance of sexual problems among the diabetics such as:

- obesity.
- hypertension.
- depression.
- lack of energy.
- mental health problems.
- Females with diabetes, show the following sexual dysfunctions:
 - Vagina dryness, as consequence of hormone changes and low flow of blood to sex organs
 - Frequent urinary infection can make diabetic women vulnerable to many sexual dysfunctions.
- Frigidity.
- Painful sexual intercourse.

Recommendations and Suggestions

1. Diabetics should get enlightened with potential risks of their disease on sexual functioning.
2. A counseling is highly needed for patients with diabetes.
3. Experimental researches should be conducted regarding the effect of diabetes on impotence.

References

1. Tamás, V., & Kempler, P. (2014). Sexual dysfunction in diabetes. *Handb Clin Neurol*, 126, 223-232. DOI: 10.1016/B978-0-444-53480-4.00017-5
2. International Diabetes Federation. (2015). Diabetes Atlas. (Fifth edition). Brussels, Belgium: *International Diabetes Federation*.
3. Campos, C. (2012). Chronic hyperglycemia and glucose toxicity: pathology and clinical sequelae. *Postgrad Med*, 124(6), 90-97. DOI: 10.3810/pgm.2012.11.2615
4. Rahman, S., Rahman, T., Ismail, A. A., & Rashid, A. R. (2007). Diabetes- associated macrovasculopathy: pathophysiology and pathogenesis. *Diabetes Obese Metab*, 9(6), 767-780. <https://doi.org/10.1111/j.1463-1326.2006.00655.x>
5. Fox, C. S., Coady, S., Sorlie, P. D., D'Agostino, R. B. Sr., Pencina, M. J., Vasan, R. S., Meigs, J. B., Levy, D., & Savage, P. J. (2007). increasing cardiovascular disease burden due to diabetes mellitus: the Framingham Heart Study. *Circulation*, 115(12), 1544-1550. DOI: 10.1161/CIRCULATIONAHA.106.658948
6. Preis, S. R., Pencina, M. J., Hwang, S. J., D'Agostino, R. B. Sr., Savage, P. J., Levy, D., & Fox, C. S. (2009). Trends in cardiovascular disease risk factors in individuals with and without diabetes mellitus in the Framingham Heart Study. *Circulation*. 120(3), 212-220. DOI: 10.1161/CIRCULATIONAHA.108.846519
7. Lu, C. C., Jiann, B. P., Sun, C. C., Lam, H. C., Chu, C. H., & Lee, J. K. (2009). Association of glycemic control with risk of erectile dysfunction in men with type 2 diabetes. *J Sex Med*, 6(6), 1719-1728. DOI: 10.1111/j.1743-6109.2009.01219.x
8. Enzlin, P., Rosen, R., Wiegel, M., Brown, J., Wessells, H., Gatcomb, P., Rutledge, B., Chan, K. L., & Cleary, P. A. (2009). DCCT/EDIC Research Group Sexual dysfunction in women with type 1 diabetes: long-term findings from the DCCT/EDIC study cohort. *Diabetes Care*. 32(5): 780-785. DOI: 10.2337/dc08-1164
9. Maseroli, E., Fanni, E., Cipriani, S., Scavello, I., Pampaloni, F., Battaglia, C., Fambrini, M., Mannucci, E., Jannini, E. A., Maggi, M., & Vignozzi, L. (2016). Cardiometabolic risk and female sexuality: focus on clitoral vascular resistance. *J Sex Med*, 13(11), 1651-1661. DOI: 10.1016/j.jsxm.2016.09.009
10. Phillips, A., & Phillips, S. (2015). Recognising female sexual dysfunction as an essential aspect of effective diabetes care. *Appl Nurs Res*, 28(3), 235-238. DOI: 10.1016/j.apnr.2015.04.007
11. Olarinoye, J., & Olarinoye, A. (2008). Determinants of sexual function among women with type 2 diabetes in a Nigerian population. *J Sex Med*, 5(4), 878-886. DOI: 10.1111/j.1743-6109.2007.00649.x
12. Giugliano, F., Maiorino, M., Bellastella, G., Gicchino, M., Giugliano, D., & Esposito, K. (2010). Determinants of erectile dysfunction in type 2 diabetes. *Int J Impot Res*, 22 (3), 204-209. DOI: 10.1038/ijir.2010.1
13. Johannes, C. B., Araujo, A. B., Feldman, H. A., Derby, C. A., Kleinman, K. P., & McKinlay, J. B. (2017). Incidence

- of erectile dysfunction in men 40 to 69 years old: longitudinal results from the Massachusetts Male Aging Study. *J Urol*, 163(2), 460-463. Retrieved from <https://pubmed.ncbi.nlm.nih.gov/10647654/>
14. Hackett, G., Krychman, M., Baldwin, D., Bennett, N., El-Zawahry, A., Graziottin, A., Lukasiewicz, M., McVary, K., Sato, Y., & Incrocci, L. (2016). Coronary heart disease, diabetes, and sexuality in men. *J Sex Med*, 13(6), 887-904. DOI: 10.1016/j.jsxm.2016.01.023
 15. Esposito, K., Maiorino, M. I., Bellastella, G., Giugliano, F., Romano, M., & Giugliano, D. (2010). Determinants of female sexual dysfunction in type 2 diabetes. *Int J Impot Res*, 22(3), 179-184. DOI: 10.1038/ijir.2010.6
 16. Seftel, A. D., Sun, P., & Swindle, R. (2004). The prevalence of hypertension, hyperlipidemia, diabetes mellitus and depression in men with erectile dysfunction. *J Urol*, 171 (6 Pt 1), 2341-2345. DOI: 10.1097/01.ju.0000125198.32936.38
 17. Kizilay, F., Gali, H. E., & Serefoglu, E. C. (2017). Diabetes and sexuality. *Sex Med Rev*, 5(1), 45-51. DOI: 10.1016/j.sxmr.2016.07.002
 18. Lewis, R. W., Fugl-Meyer, K. S., Corona, G., Hayes, R. D., Laumann, E. O., Moreira, E. D. Jr., Rellini, A. H., & Segraves, T. (2010). Definitions/epidemiology/risk factors for sexual dysfunction. *J Sex Med*, 7(4 Pt 2), 1598-1607. DOI: 10.1111/j.1743-6109.2010.01778.x
 19. Miner, M., Esposito, K., Guay, A., Montorsi, P., & Goldstein, I. (2012). Cardio metabolic risk and female sexual health: the Princeton III summary. *J Sex Med*, 9(3), 641-651. quiz 652. DOI: 10.1111/j.1743-6109.2012.02649.x
 20. Esposito, K., Ciotola, M., Maiorino, M. I., Giugliano, F., Autorino, R., De Sio, M., Cozzolino, D., Saccomanno, F., & Giugliano, D. (2009). Hyperlipidemia and sexual function in premenopausal women. *J Sex Med*, 6(6): 1696-1703. DOI: 10.1111/j.1743-6109.2009.01284.x
 21. Esposito, K., Ciotola, M., Marfella, R., Di Tommaso, D., Cobellis, L., & Giugliano, D. (2005). Sexual dysfunction in women with the metabolic syndrome. *Diabetes Care*, 28(3), 756. Retrieved from <https://agris.fao.org/agris-search/search.do?recordID=US201300989451>
 22. Isidori, A. M., Buvat, J., Corona, G., Goldstein, I., Jannini, E. A., Lenzi, A., Porst, H., Salonia, A., Traish, A. M., & Maggi, M. (2017). A critical analysis of the role of testosterone in erectile function: from pathophysiology to treatment – a systematic review. *Eur Urol*, 65(1), 99-112. DOI: 10.1016/j.eururo.2013.08.048
 23. Bellastella, G., Maiorino, M. I., Olita, L., De Bellis, A., Giugliano, D., & Esposito, K. (2013). Anti-pituitary antibodies and hypogonadotropic hypogonadism in type 2 diabetes: in search of a role. *Diabetes Care*, 36(8), e116-e117. DOI: 10.2337/dc13-0637
 24. Meeking, D. R., Fosbury, J. A., & Cummings, M. H. (2016). Sexual dysfunction and sexual health concerns in women with diabetes. *Practical Diabetes*, 30(8), 327-331. <https://doi.org/10.1002/pdi.1805>
 25. Pohnholzer, A., Temml, C., Mock, K., Marszalek, M., Obermayr, R., & Madersbacher, S. (2005). Prevalence and risk factors for erectile dysfunction in 2869 men using a validated questionnaire. *Eur Urol*, 47(1), 80-85. discussion 85-86. DOI: 10.1016/j.eururo.2004.08.017
 26. Go, A. S., Mozaffarian, D., Roger, V. L., Benjamin, E. J., Berry, J. D., Borden, W. B., Bravata, D. M., Dai, S., Ford, E. S., Fox, C. S., Franco, S., Fullerton, H. J., Gillespie, C., Hailpern, S. M., Heit, J. A., Howard, V. J., Huffman, M. D., Kissela, B. M., Kittner, S. J., ... Turner, M. B. (2013). American Heart Association Statistics Committee and Stroke Statistics Subcommittee Heart disease and stroke statistics – 2013 update: a report from the American Heart Association. *Circulation*, 127(1), e6-e245. DOI: 10.1161/CIR.0b013e31828124ad
 27. Giuliano, F. A., Leriche, A., Jaudinot, E. O., & de Gendre, A. S. (2004). Prevalence of erectile dysfunction among 7689 patients with diabetes or hypertension, or both. *Urology*, 64(6), 1196-1201. DOI: 10.1016/j.urology.2004.08.059
 28. Maria Ida, Maiorino., Giuseppe, Bellastella., & Katherine, Esposito., (2014). Diabetes and sexual dysfunction: current perspectives *Diabetes Metab Syndr Obes*, 7, 95-105. DOI: 10.2147/DMSO.S36455
 29. Demir, O., Akgul, K., Akar, Z., Cakmak, O., Ozdemir, I., Bolukbasi, A., Can, E., & Gumus, B. H. (2009). Association between severity of lower urinary tract symptoms, erectile dysfunction and metabolic syndrome. *Aging Male*, 12(1), 29-34. DOI: 10.1080/13685530902777425
 30. Turek, S. J., Hastings, S. M., Sun, J. K., King, G. L., & Keenan, H. A. (2013). Sexual dysfunction as a marker of cardiovascular disease in males with 50 or more years of type 1 diabetes. *Diabetes Care*, 36(10), 3222-3226. DOI: 10.2337/dc13-0294
 31. Dong, J. Y., Zhang, Y. H., & Qin, L. Q. (2011). Erectile dysfunction and risk of cardiovascular disease: meta-analysis of prospective cohort studies. *J Am Coll Cardiol*, 58(13), 1378-1385. DOI: 10.1016/j.jacc.2011.06.024
 32. Araujo, A. B., Travison, T. G., Ganz, P., Chiu, G. R., Kupelian, V., Rosen, R. C., Hall, S. A., & McKinlay, J. B. (2009). Erectile dysfunction and mortality. *J Sex Med*, 6(9), 2445-2454. DOI: 10.1111/j.1743-6109.2009.01354.x
 33. Thorve, V. S., Kshirsagar, A. D., Vyawahare, N. S., Joshi, V. S., Ingale, K. G., & Mohite, R. J. (2011). Diabetes-induced erectile dysfunction: epidemiology, pathophysiology and management. *J Diabetes Complications*, 25(2), 129-136. DOI: 10.1016/j.jdiacomp.2010.03.003
 34. Corona, G., Giorda, C. B., Cucinotta, D., Guida, P., & Nada, E. (2013). Gruppo di studio SUBITO-DE The SUBITO-DE study: sexual dysfunction in newly diagnosed type 2 diabetes male patients. *J Endocrinol Invest*, 36(10), 864-868. DOI: 10.3275/8969
 35. Chew, S. K. h., Taouk, Y., Xie, J., Nicolaou, T. E., Wang, J. J., Wong, T. Y., & Lamoureux, E. L. (2014). Relationship between diabetic retinopathy, diabetic macular oedema and erectile dysfunction in type 2 diabetics. *Clin Experiment Ophthalmol*, 41(7), 683-689. DOI: 10.1111/ceo.12099
 36. Al-Hunayan, A., Al-Mutar, M., Kehinde, E. O., Thalib, L., & Al-Ghorory, M. (2007). The prevalence and predictors of erectile dysfunction in men with newly diagnosed with type 2 diabetes mellitus. *BJU Int*, 99(1), 130-134.

DOI: 10.1111/j.1464-410X.2006.06550.x

37. Rosen, R. C., Wing, R. R., Schneider, S., Wadden, T. A., Foster, G. D., West, D. S., Kitabchi, A. E., Brancati, F. L., Maschak-Carey, B. J., Bahnson, J. L., Lewis, C. E., & Gendrano, I. N. (2009). Erectile dysfunction in type 2 diabetic men: relationship to exercise fitness and cardiovascular risk factors in the Look AHEAD trial. *J Sex Med*, 6(5), 1414–1422.
DOI: 10.1111/j.1743-6109.2008.01209.x
38. Corona, G., Giorda, C. B., Cucinotta, D., Guida, P., & Nada, E., Gruppo di studio SUBITO-DE The SUBITO-DE study: sexual dysfunction in newly diagnosed type 2 diabetes male patients. *J Endocrinol Invest*, 36(10), 864–868. DOI: 10.3275/8969
39. Chew, S. K. h., Taouk, Y., Xie, J., Nicolaou, T. E., Wang, J. J., Wong, T. Y., & Lamoureux, E. L. (2013). Relationship between diabetic retinopathy, diabetic macular oedema and erectile dysfunction in type 2 diabetics. *Clinical Experiment Ophthalmology*, 41(7), 683–689.
DOI: 10.1111/ceo.12099
40. Heruti, R. J., Uri, I., Arbel, Y., Swartzon, M., Galor, S., & Justo, D. (2007). Erectile dysfunction severity might be associated with poor cardiovascular prognosis in diabetic men. *J Sex Med*, 4(2), 465–471.
DOI: 10.1111/j.1743-6109.2006.00420.x
41. Chew, K. K., Bremner, A., Jamrozik, K., Earle, C., & Stuckey, B. (2008). Male erectile dysfunction and cardiovascular disease: is there an intimate nexus? *J Sex Med*, 5(4), 928–934. <https://doi.org/10.1111/j.1743-6109.2007.00714.x>
42. Foresta, C., Caretta, N., Corona, G., Fabbri, A., Francavilla, S., Jannini, E., Maggi, M., Bettocchi, C., & Lenzi, A. (2009). Clinical and metabolic evaluation of subjects with erectile dysfunction: a review with a proposal flowchart. *Int J Androl*, 32(3), 198–211. DOI:10.1111/j.1365-2605.2008.00932.x
43. Hackett, G., Krychman, M., Baldwin, D., Baldwin, D., El-Zawahry, A., Graziottin, A., Lukasiewicz, M., McVary, K., Sato, Y., & Incrocci, L. (2016). Coronary heart disease, diabetes, and sexuality in men. *J Sex Med*, 13(6), 887-904.
DOI: 10.1016/j.jsxm.2016.01.023
44. Bal, M. D., Yılmaz, S. D., Celik, S. G., Dinçaç, N., Beji, N. K., & Yalçın, O. (2015). Does the diabetes of type 2 affect the sexual functions of women? *J Sex Marital Ther*, 41(1), 107-13. DOI: 10.1080/0092623X.2013.842193
45. Thorve, V. S., Kshirsagar, A. D., Vyawahare, N. S., Joshi, V. S., Ingale, K. G., & Mohite, R. J. (2011). Diabetes-induced erectile dysfunction: epidemiology, pathophysiology and management. *J Diabetes Complications*, 25(2), 129-136.
DOI: 10.1016/j.jdiacomp.2010.03.003

Copyright: ©2022 Ibrahim Abdelrahim Ibrahim Humaida. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.