

Medication Adherence: Fact or Fictions?

Abdul Kader Mohiuddin

*Alumni, Faculty of Pharmacy, Dhaka University****Correspondence author****Abdul Kader Mohiuddin**

Alumni

Faculty of Pharmacy

Dhaka University

Submitted : 15 Nov 2022 ; Published : 13 Dec 2022

Citation : Abdul Kader Mohiuddin, Medication Adherence: Fact or Fictions?. J Nurs Care Repo; 3(3): 1-3.**Keywords:** Patient compliance, Healthcare denials, Medication adherence, Elderly patient care, Treatment failure.

Proper use of medicine or taking medicine in correct order is essential to cure any disease. According to the WHO, lack of adherence to treatment regimens leads to major problems among patients, mostly with chronic illnesses. "Right administration" depends on at least 5 right factors--right patient, right drug, right time, right dose and right route (Grissinger, 2010). "Medicines won't work if you don't take it right"--This simple fact is not understood by most people in the world, as a result still more than half of the patients with chronic diseases in the developed world do not take their medicine correctly--says WHO (Brown & Bussell, 2011).

Patients with chronic diseases may find it particularly difficult to adhere because they frequently need to take their medications for an extended period of time, sometimes for the remainder of their lives. There are several reasons why patients may find it challenging to adhere to treatment regimens, and CDC estimates that medication non-compliance leads to 30 to 50% of chronic disease treatment failures. Poor adherence may cause treatment outcomes to not be achieved, symptoms to worsen, and one's health to deteriorate (Center for Drug Evaluation and Research, (n.d)).

In the UK, up to 50% of medicines are not taken as intended and 60% of NHS patients failed to receive the right treatment within 18 weeks (Barnett, 2014 ; Campbell et al., 2022 ; Andrews & Luke, 2022). According to a New Zealand-based Journal medication non-adherence alone accounts for at least 10% of hospitalizations, \$300 billion in annual medical costs, and more than 1,25, 000 deaths in the US alone (Cutler et al., 2019). A recent Canadian study found that 30% of patients stop taking their medication before it is instructed, and one in four do not fill their prescription or take less than prescribed (Bonsu et al., 2022). Medication non-adherence contributes more than half of the \$500 billion total avoidable costs attributed to suboptimal medicine use globally each year (Furniss et al., 2014).

A strange similarity can be found in under-developed, developing countries and the so-called developed world in the West or the Middle-East when it comes to not taking medicine

properly. More than half of the antibiotics worldwide are sold without a prescription and CDC stated, 30-50% of antibiotics prescribed in hospitals are inappropriate or unnecessary (Bahta et al., 2020; Improve Antibiotic Use, 2022). A recent study published by The Lancet, funded by the Bill & Melinda Gates Foundation, Wellcome Trust, states that nearly 5 million deaths worldwide in 2019 were related to bacterial resistance, which is expected to double by 2050 (Antimicrobial Resistance Collaborators, 2022).

Globally, NSAIDs are responsible for at least 650,000 hospitalizations and 165,000 deaths annually (Kasciuškevičiūtė et al., 2018). Overuse of this class of drugs can cause kidney injury, and their side effects can be 3 to 4 times higher in kidney compromised patients (Lucas et al., 2019). Many studies have reported widespread misuse of these drugs in Dengue, Chikungunya and Covid-19 patients. Especially in Dengue or Covid-19 patients, it is more important to maintain the hydration level of the body than to bring down the fever with the pain killers. In children, the use of excess Paracetamol syrup or suppositories may cause stomach irritation, which hampers digestion and led to vomiting and ended up with hospitalization. Most hospitalizations or ICU admissions among those patients could be prevented, with few exceptions, simply by preventing dehydration at home with saline and fruit juice or simply by drinking more water.

More or less 40% of Covid-19 patients report sleep disturbances--use of Benzodiazepines in Covid-19 patients increases the incidence of delirium, depresses the system in patients with compromised respiratory functions, and contraindicated with some anti-viral medications (Jahrami et al., 2021 ; Ostuzzi et al., 2020).

Around 300 metric tons of morphine-type painkillers are used worldwide each year, less than 1% of which distributed to low-and-middle income countries, says the American Journal of Public Health (Bhadelia et al., 2019). So their misuse and related side effects are also retained by the developed world.

According to a WHO report, only half of patients in developed countries adhere to treatment guidelines for chronic diseases, which is much less in developing countries (Chauke et al., 2022). Several studies among diabetic patients in South Asian countries have shown that nearly half of patients do not adhere to their prescribed medication and are at risk of acute and long-term complications, resulting in increased hospitalization rates and medical costs (Chong et al., 2014; Sohal et al., 2015).

Three-quarters of elderly patients worldwide are unable to adhere to appropriate long-term treatment regimens—due to multiple physical complications and additional medication burden (Félix & Henriques, 2021). Elderly patients taking at least 5 medications are at increased risk of mild cognitive impairment, dementia, falls, frailty, disability, and mortality, while ADEs are estimated to be 5% to 28% of acute geriatric medical admissions (Chippa & Roy, 2022 ; Varghese et al., 2022).

A recent study by the American Heart Association revealed that patients with high blood pressure do not follow treatment guidelines because of--(1) suboptimal dosing or prescribing the wrong medication (2) lack of insurance or lack of health care access and (3) patient failure to comply prescribed medication or other lifestyle guidelines (Choudhry et al., 2022).

“Medical costs are barriers to adherence to proper clinical guidelines for long-term diseases in poor countries”-- although discussed in many forums but forgetfulness, confusion about the duration required for medication use and mistrust about the overall efficacy of medication are among the reasons for non-adherence to diabetes management protocols in Middle Eastern countries (Alsairafi et al., 2016).

Epilepsy is still seen in most countries of the world as an evil spirit --although two-thirds of patients can become seizure-free with adequate treatment, poor adherence to proper guidelines is a major problem for effective recovery (Lossius et al., 2017).

Several identified reasons for non-adherence to treatment guidelines for chronic diseases:

1. **Patient’s socio-economic status:** low health literacy, lack of family or social support network, unstable living or homelessness, financial insecurity
2. **Treatment-related:** complexity and duration of treatment procedures, frequent changes in medication regimen, lack of immediate results, real or perceived unpleasant side effects, interference with lifestyle
3. **Health system-related:** high treatment costs, limited health system for patient education and follow-up, doctor-patient relationship, patient trust in health care, long waits, lack of patient information materials
4. **Patient-related:** visual-hearing and cognitive impairment, mobility and dexterity, psychological and behavioral factors, perceived risk of disease susceptibility, superstitions and stigmatization by disease, etc. (Jin et al., 2008)

Finally, it can be said that patients’ knowledge and perception of the disease is the main driving force in determining their adherence to the treatment regimen. Health care providers should explore providing more effective health-education to identify patients’ attitudes toward disease, trust in medications, psychological stressors, and increase medication adherence.

Acknowledgement

I’m thankful to Khulud Alosaimi, Department of Pharmacology and Toxicology, Faculty of Medicine, Umm Al-Qura University, Makkah for his valuable time to audit my paper and for his thoughtful suggestions. I’m also grateful to seminar library of Faculty of Pharmacy, University of Dhaka and BANSDOC Library, Bangladesh for providing me books, journal and newsletters.

Abbreviations

Adverse drug events (ADEs)
Non-steroidal anti-inflammatory drugs (NSAIDs)
The Centers for Disease Control and Prevention (CDC)
World Health Organization (WHO)

Financial Disclosure or Funding: N/A

Conflict of Interest: The author declares that he has no competing interests.

Informed Consent: N/A

References

1. Grissinger, M. (2010). The Five Rights: A Destination Without a Map. *P T*, 35(10), 542. Retrieved from https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2957754/pdf/ptj35_10p542.pdf
2. Brown, M. T., & Bussell, J. K. (2011). Medication adherence: WHO cares? *Mayo Clin Proc*, 86(4), 304-14. DOI: 10.4065/mcp.2010.0575
3. Center for Drug Evaluation and Research. (n.d.). Why you need to take your medications as prescribed or instructed. U.S. *Food and Drug Administration*. November 5, 2022, Retrieved from <https://www.fda.gov/drugs/special-features/why-you-need-to-take-your-medications-prescribed-or-instructed>
4. Barnett, N. L. (2013). Medication adherence: where are we now? A UK perspective. *European Journal of Hospital Pharmacy*, 21(3), 181-184. DOI:10.1136/ejhp-2013-000373
5. Campbell, Denis, and Pamela Duncan. (2022). “Record 6.8m People Waiting for Hospital Treatment in England.” *The Guardian*, 8 Sept. 2022, www.theguardian.com/society/2022/sep/08/waiting-lists-for-routine-hospital-treatment-in-england-break-record.
6. Andrews, Luke. (2022). “60% of Patients Waiting 18 Wks for Treatment at Worst-performing Trust.” *Mail Online*, 7 Mar. 2022, www.dailymail.co.uk/news/article-10577387/60-NHS-patients-waited-18-weeks-treatment-worst-performing-hospital-trust.html.
7. Cutler, R. L., Torres-Robles, A., Wiecek, E., Drake, B., Van der Linden, N., Benrimoj, S. I., & Garcia-Cardenas, V. (2019). Pharmacist-led medication non-adherence

- intervention: reducing the economic burden placed on the Australian health care system. *Patient Prefer Adherence*, 13, 853-862. <https://doi.org/10.2147/PPA.S191482>
8. Bonsu, K. O., Young, S., Lee, T., Nguyen, H., & Chitsike, R. S. (2022). Adherence to Antithrombotic Therapy for Patients Attending a Multidisciplinary Thrombosis Service in Canada - A Cross-Sectional Survey. *Patient Prefer Adherence*, 26(16), 1771-1780. DOI: 10.2147/PPA.S367105
 9. Furniss, D., Barber, N., Lyons, I., Eliasson, L., & Blandford, A. (2014). Unintentional non-adherence: can a spoon full of resilience help the medicine go down? *BMJ Qual Saf*, 23(2), 95-8. DOI: 10.1136/bmjqs-2013-002276
 10. Bahta, M., Tesfamariam, S., Weldemariam, D. G., Yemane, H., Tesfamariam, E. H., Alem, T., & Russom, M. (2020). Dispensing of antibiotics without prescription and associated factors in drug retail outlets of Eritrea: A simulated client method. *PLoS One*, 15(1), e0228013. <https://doi.org/10.1371/journal.pone.0228013>
 11. "Improve Antibiotic Use. (2022)." Centers for Disease Control and Prevention, Centers for Disease Control and Prevention." Retrieved from <https://www.cdc.gov/sixteenths/hai/index.htm>.
 12. Antimicrobial Resistance Collaborators. (2022). Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *Lancet*, 399(10325), 629-655. DOI: 10.1016/S0140-6736(21)02724-0
 13. Kasciūškevičiūtė, S., Gumbrevičius, G., Vendzelytė, A., Ščiupokas, A., Petrikonis, K., & Kaduševičius, E. (2018). Impact of the World Health Organization Pain Treatment Guidelines and the European Medicines Agency Safety Recommendations on Nonsteroidal Anti-Inflammatory Drug Use in Lithuania: An Observational Study. *Medicina (Kaunas)*, 54(2), 30. DOI: 10.3390/medicina54020030
 14. Lucas, G. N. C., Leitão, A. C. C., Alencar, R. L., Xavier, R. M. F., Daher, E. F., Silva Junior, G. B. D. (2019). Pathophysiological aspects of nephropathy caused by non-steroidal anti-inflammatory drugs. *J Bras Nefrol*, 41(1), 124-130. DOI: 10.1590/2175-8239-JBN-2018-0107
 15. Jahrami, H., BaHammam, A. S., Bragazzi, N. L., Saif, Z., Faris, M., & Vitiello, M. V. (2021). Sleep problems during the COVID-19 pandemic by population: a systematic review and meta-analysis. *J Clin Sleep Med*, 17(2), 299-313. DOI: 10.5664/jcsm.8930
 16. Ostuzzi, G., Papola, D., Gastaldon, C., Schoretsanitis, G., Bertolini, F., Amaddeo, F., Cuomo, A., Emsley, R., Fagiolini, A., Imperadore, G., Kishimoto, T., Michencigh, G., Nosé, M., Purgato, M., Dursun, S., Stubbs, B., Taylor, D., Thornicroft, G., Ward, P. B., Hiemke, C., Correll, C. U., & Barbui, C. (2020). Safety of psychotropic medications in people with COVID-19: evidence review and practical recommendations. *BMC Med*, 18(1), 215. <https://doi.org/10.1186/s12916-020-01685-9>
 17. Bhadelia, A., De Lima, L., Arreola-Ornelas, H., Kwete, X. J., Rodriguez, N. M., & Knaul, F. M. (2019). Solving the Global Crisis in Access to Pain Relief: Lessons From Country Actions. *Am J Public Health*, 109(1), 58-60. DOI: 10.2105/AJPH.2018.304769
 18. Chauke, G. D., Nakwafila, O., Chibi, B., Sartorius, B., Mashamba-Thompson, T. (2022). Factors influencing poor medication adherence amongst patients with chronic disease in low-and-middle-income countries: A systematic scoping review. *Heliyon*, 8(6), e09716. DOI: 10.1016/j.heliyon.2022.e09716
 19. Chong, E., Wang, H., King-Shier, K. M., Quan, H., Rabi, D. M., & Khan N. A. (2014). Prescribing patterns and adherence to medication among South-Asian, Chinese and white people with Type 2 diabetes mellitus: a population-based cohort study. *Diabet Med*, 31(12), 2014. DOI: 10.1111/dme.12559
 20. Sohal, T., Sohal, P., King-Shier, K. M., & Khan, N. A. (2015). Barriers and Facilitators for Type-2 Diabetes Management in South Asians: A Systematic Review. *PLoS One*, 10(9), e0136202. DOI: 10.1371/journal.pone.0136202
 21. Félix, I. B., Henriques, A. (2021). Medication adherence and related determinants in older people with multimorbidity: a cross-sectional study. *Nurs Forum*, 56(4), 834-843. DOI: 10.1111/nuf.12619
 22. Chippa, V., Roy, K. (2022). Geriatric Cognitive Decline and Polypharmacy. [Updated 2022 Apr 1]. In: StatPearls [Internet]. *Treasure Island (FL): StatPearls Publishing*, Retrieved from <https://www.ncbi.nlm.nih.gov/books/NBK574575/>
 23. Varghese, D., Ishida, C., & Haseer Koya, H. (2022). Polypharmacy. [Updated 2022 Sep 9]. In: StatPearls [Internet]. *Treasure Island (FL): StatPearls Publishing*, Retrieved from <https://www.ncbi.nlm.nih.gov/books/NBK532953/>
 24. Choudhry, N. K., Kronish, I. M., Vongpatanasin, W., Ferdinand, K. C., Pavlik, V. N., Egan, B. M., Schoenthaler, A., Houston Miller, N., & Hyman, D. J. (2022). American Heart Association Council on Hypertension; Council on Cardiovascular and Stroke Nursing; and Council on Clinical Cardiology. Medication Adherence and Blood Pressure Control: A Scientific Statement From the American Heart Association. *Hypertension*, 79(1), e1-e14. DOI: 10.1161/HYP.0000000000000203
 25. Alsairafi, Z. K., Taylor, K. M., Smith, F. J., Alattar, A. T. (2016). Patients' management of type 2 diabetes in Middle Eastern countries: review of studies. *Patient Prefer Adherence*, 10(10), 1051-62. DOI: 10.2147/PPA.S104335
 26. Lossius, M. I., Alfstad, K. Å., Aaberg, K. M., & Nakken, K. O. (2017). Seponering av Antiepileptika Ved anfallsfrihet – når og hvordan? *Tidsskrift for Den Norske Legeforening*, 137(6), 451-454. <https://doi.org/10.4045/tidsskr.16.0957>
 27. Jin, J., Sklar, G. E., Min Sen Oh, V., & Chuen Li, S. (2008). Factors affecting therapeutic compliance: A review from the patient's perspective. *Ther Clin Risk Manag*, 4(1), 269-86. DOI : 10.2147/tcrm.s1458

Copyright: ©2022 Abdul Kader Mohiuddin. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.